

Kool Kids Club

Infectious and Communicable Diseases

Our Club is committed to the health and safety of all children and Staff, who play, learn and work here. As such, it will sometimes be necessary to require a poorly child to be collected early from a session or be kept at home while they get better. In such cases, the provisions of the Health, Illness and Emergency policy will be implemented.

In accordance with the procedures set out in the Health, Illness and Emergency policy, parents/carers will be notified immediately if their child has become ill and needs to go home. Poorly children will be comforted, kept safe and under close supervision until they are collected.

If a child has had to go home prematurely due to illness, they should remain at home until they are better for at least 48 hours, or according to the times set out in the table below. If a member of staff becomes ill at work, similar restrictions on their return will apply.

If a child or member of staff becomes ill outside Club hours, they should notify the Club as soon as possible. The minimum exclusion periods outlined in the table on the next page will then come into operation.

If any infectious or communicable disease is detected on the Club's premises, the Club will inform parents/carers personally in writing as soon as possible. The Club is committed to sharing as much information as possible about the source of the disease and the steps being taken to remove it. Ofsted will also be informed of any infectious or communicable diseases discovered on the Club's premises.

Head lice

When a case of head lice is discovered at the Club, the situation will be handled carefully, sensitively and safely. The child concerned will not be isolated from other children, and there is no need for them to be excluded from activities or sessions at the Club.

When the child concerned is collected, their parent/carer will be informed in a sensitive manner. Other parents/carers will be informed as quickly as possible in writing, including advice and guidance on treating head lice.

Staff should check themselves regularly for lice and treat whenever necessary.

Chicken Pox S days from onset of rash and scabs have formed. Respiratory infections including coronavirus (COVID-19) Children and young people should not attend if they have a high temperature and are unwell Children with mild symptoms such as runny nose, and headache who are otherwise well can continue to attend settings/school. German Measles (Rubella) German Measles (Rubella) S days from onset of rash (Over the Rainbow must notify staff/mothers who may be pregnant) None. Chemical treatment is recommended only in cases where live lice have been seen. Combing may be more effective. Hand, Foot, and Mouth None once child is well Impetigo Until lesions are crusted or healed Influenza None once child is well Measles Four days from onset of rash (Kool Kids must notify staff/mothers who may be pregnant) Meningococcal Meningitis/Septicaemia Will give specific advice on any action needed. Siblings need not be excluded Meningitis None once child is well infection risk is minimal Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea 48 hours after the last bout of either. None. (Kool Kids must notify staff/mothers who may be pregnant) None. (Kool Kids must notify staff/mothers who may be pregnant) None. (Kool Kids must notify staff/mothers who may be pregnant) None. (Kool Kids must notify staff/mothers who may be pregnant) None once child is well S days from commencement of treatment			
Respiratory infections including coronavirus (COVID-19) Children and young people should not attend if they have a high temperature and are unwell Children with mild symptoms such as runny nose, and headache who are otherwise well can continue to attend settings/school. German Measles (Rubella) German Measles (Rubella) S days from onset of rash (Over the Rainbow must notify staff/mothers who may be pregnant) Head Lice None. Chemical treatment is recommended only in cases where live lice have been seen. Combing may be more effective. None once child is well Impetigo Until lesions are crusted or healed Influenza None once child is well Measles Four days from onset of rash (Kool Kids must notify staff/mothers who may be pregnant) Meningococcal Meningitis/Septicaemia Meningitis/Septicaemia Meningitis None once child is well infection risk is minimal Mumps Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea Slapped Cheek Syndrome (Parvovirus B19) (Kool Kids must notify staff/mothers who may be pregnant) None. (Kool Kids must notify staff/mothers who may be pregnant)	DISEASE/ILLNESS	EXCLUSION PERIOD	
they have a high temperature and are unwell Children with mild symptoms such as runny nose, and headache who are otherwise well can continue to attend settings/school. German Measles (Rubella) S days from onset of rash (Over the Rainbow must notify staff/mothers who may be pregnant) Head Lice None. Chemical treatment is recommended only in cases where live lice have been seen. Combing may be more effective. Hand, Foot, and Mouth None once child is well Impetigo Until lesions are crusted or healed Influenza None once child is well Four days from onset of rash (Kool Kids must notify staff/mothers who may be pregnant) Meningococcal Meningitis/Septicaemia Siblings need not be excluded Meningitis None once child is well infection risk is minimal Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea Slapped Cheek Syndrome (Parvovirus B19) (Kool Kids must notify staff/mothers who may be pregnant) None. (Kool Kids must notify staff/mothers who may be pregnant) Kool Kids must notify staff/mothers who may be pregnant) None.	Chicken Pox	5 days from onset of rash and scabs have formed.	
Children with mild symptoms such as runny nose, and headache who are otherwise well can continue to attend settings/school. German Measles (Rubella) 5 days from onset of rash (Over the Rainbow must notify staff/mothers who may be pregnant) Head Lice None. Chemical treatment is recommended only in cases where live lice have been seen. Combing may be more effective. Hand, Foot, and Mouth None once child is well Impetigo Until lesions are crusted or healed Influenza None once child is well Measles Four days from onset of rash (Kool Kids must notify staff/mothers who may be pregnant) Meningococcal Meningitis/Septicaemia Mil give specific advice on any action needed. Siblings need not be excluded Meningitis None once child is well infection risk is minimal Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea Slapped Cheek Syndrome (Parvovirus B19) (Kool Kids must notify staff/mothers who may be pregnant) None. (Kool Kids must notify staff/mothers who may be pregnant) None once child is well	Respiratory infections including	Children and young people should not attend if	
and headache who are otherwise well can continue to attend settings/school. German Measles (Rubella) 5 days from onset of rash (Over the Rainbow must notify staff/mothers who may be pregnant) Head Lice None. Chemical treatment is recommended only in cases where live lice have been seen. Combing may be more effective. Hand, Foot, and Mouth None once child is well Impetigo Until lesions are crusted or healed Influenza None once child is well Measles Four days from onset of rash (Kool Kids must notify staff/mothers who may be pregnant) Meningococcal Meningitis/Septicaemia Mil give specific advice on any action needed. Siblings need not be excluded Meningitis None once child is well infection risk is minimal Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea Slapped Cheek Syndrome (Parvovirus B19) (Kool Kids must notify staff/mothers who may be pregnant) None once child is well	coronavirus (COVID-19)	they have a high temperature and are unwell	
to attend settings/school. German Measles (Rubella) 5 days from onset of rash (Over the Rainbow must notify staff/mothers who may be pregnant) Head Lice None. Chemical treatment is recommended only in cases where live lice have been seen. Combing may be more effective. Hand, Foot, and Mouth None once child is well Impetigo Until lesions are crusted or healed Influenza None once child is well Four days from onset of rash (Kool Kids must notify staff/mothers who may be pregnant) Meningococcal The consultant in communicable disease control will give specific advice on any action needed. Siblings need not be excluded Meningitis None once child is well infection risk is minimal Mumps Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea Slapped Cheek Syndrome (Parvovirus B19) (Kool Kids must notify staff/mothers who may be pregnant) None. (Kool Kids must notify staff/mothers who may be pregnant) None once child is well			
German Measles (Rubella) 5 days from onset of rash (Over the Rainbow must notify staff/mothers who may be pregnant) Head Lice None. Chemical treatment is recommended only in cases where live lice have been seen. Combing may be more effective. Hand, Foot, and Mouth None once child is well Impetigo Until lesions are crusted or healed Influenza None once child is well Four days from onset of rash (Kool Kids must notify staff/mothers who may be pregnant) Meningococcal Meningitis/Septicaemia Will give specific advice on any action needed. Siblings need not be excluded Meningitis None once child is well infection risk is minimal Mumps Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea 48 hours after the last bout of either. None. (Kool Kids must notify staff/mothers who may be pregnant) None once child is well			
(Over the Rainbow must notify staff/mothers who may be pregnant) Head Lice None. Chemical treatment is recommended only in cases where live lice have been seen. Combing may be more effective. Hand, Foot, and Mouth None once child is well Impetigo Until lesions are crusted or healed Influenza None once child is well Measles Four days from onset of rash (Kool Kids must notify staff/mothers who may be pregnant) Meningococcal Meningitis/Septicaemia Will give specific advice on any action needed. Siblings need not be excluded Meningitis None once child is well infection risk is minimal Mumps Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea 48 hours after the last bout of either. None. (Kool Klds must notify staff/mothers who may be pregnant) Tonsillitis None once child is well			
may be pregnant) Head Lice None. Chemical treatment is recommended only in cases where live lice have been seen. Combing may be more effective. Hand, Foot, and Mouth None once child is well Impetigo Until lesions are crusted or healed Influenza None once child is well Measles Four days from onset of rash (Kool Kids must notify staff/mothers who may be pregnant) Meningococcal The consultant in communicable disease control will give specific advice on any action needed. Siblings need not be excluded Meningitis None once child is well infection risk is minimal Mumps Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea 48 hours after the last bout of either. None. (Kool Klds must notify staff/mothers who may be pregnant) Tonsillitis None once child is well	German Measles (Rubella)	· ·	
Head Lice None. Chemical treatment is recommended only in cases where live lice have been seen. Combing may be more effective. Hand, Foot, and Mouth None once child is well Impetigo Until lesions are crusted or healed Influenza None once child is well Measles Four days from onset of rash (Kool Kids must notify staff/mothers who may be pregnant) Meningococcal The consultant in communicable disease control will give specific advice on any action needed. Siblings need not be excluded Meningitis None once child is well infection risk is minimal Mumps Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea 48 hours after the last bout of either. Slapped Cheek Syndrome (Parvovirus None. (Kool Klds must notify staff/mothers who may be pregnant) Tonsillitis None once child is well			
cases where live lice have been seen. Combing may be more effective. Hand, Foot, and Mouth Impetigo Until lesions are crusted or healed Influenza None once child is well Measles Four days from onset of rash (Kool Kids must notify staff/mothers who may be pregnant) Meningococcal Meningitis/Septicaemia Will give specific advice on any action needed. Siblings need not be excluded Meningitis None once child is well infection risk is minimal Mumps Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea Slapped Cheek Syndrome (Parvovirus B19) (Kool Klds must notify staff/mothers who may be pregnant) Tonsillitis None once child is well			
be more effective. Hand, Foot, and Mouth None once child is well Impetigo Until lesions are crusted or healed Influenza None once child is well Measles Four days from onset of rash (Kool Kids must notify staff/mothers who may be pregnant) Meningococcal Meningitis/Septicaemia Will give specific advice on any action needed. Siblings need not be excluded Meningitis None once child is well infection risk is minimal Mumps Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea Slapped Cheek Syndrome (Parvovirus B19) (Kool Klds must notify staff/mothers who may be Fifth Disease Tonsillitis None once child is well	Head Lice	•	
Hand, Foot, and Mouth Impetigo Until lesions are crusted or healed Influenza None once child is well Measles Four days from onset of rash (Kool Kids must notify staff/mothers who may be pregnant) Meningococcal Meningitis/Septicaemia Will give specific advice on any action needed. Siblings need not be excluded Meningitis None once child is well infection risk is minimal Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea Slapped Cheek Syndrome (Parvovirus B19) (Kool Kids must notify staff/mothers who may be pregnant) Tonsillitis None once child is well			
Impetigo Until lesions are crusted or healed Influenza None once child is well Measles Four days from onset of rash (Kool Kids must notify staff/mothers who may be pregnant) Meningococcal The consultant in communicable disease control will give specific advice on any action needed. Siblings need not be excluded Meningitis None once child is well infection risk is minimal Mumps Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea 48 hours after the last bout of either. Slapped Cheek Syndrome (Parvovirus None. (Kool Kids must notify staff/mothers who may be pregnant) Tonsillitis None once child is well			
Influenza Measles Four days from onset of rash (Kool Kids must notify staff/mothers who may be pregnant) Meningococcal Meningitis/Septicaemia Meningitis Meningitis Meningitis Meningitis None once child is well infection risk is minimal Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea Slapped Cheek Syndrome (Parvovirus B19) (Kool Klds must notify staff/mothers who may be Fifth Disease Tonsillitis None once child is well	,		
Measles Four days from onset of rash (Kool Kids must notify staff/mothers who may be pregnant) Meningococcal Meningitis/Septicaemia Meningitis Meningitis Mone once child is well infection risk is minimal Mumps Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea 48 hours after the last bout of either. Slapped Cheek Syndrome (Parvovirus B19) (Kool Klds must notify staff/mothers who may be pregnant) Tonsillitis None once child is well			
(Kool Kids must notify staff/mothers who may be pregnant) Meningococcal The consultant in communicable disease control will give specific advice on any action needed. Siblings need not be excluded Meningitis None once child is well infection risk is minimal Mumps Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea 48 hours after the last bout of either. Slapped Cheek Syndrome (Parvovirus None. (Kool Klds must notify staff/mothers who may be pregnant) Tonsillitis None once child is well			
Meningococcal Meningitis/Septicaemia Meningitis/Septicaemia Meningitis Meningitis Meningitis Meningitis Meningitis Meningitis None once child is well infection risk is minimal Mumps Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea 48 hours after the last bout of either. Slapped Cheek Syndrome (Parvovirus B19) (Kool Klds must notify staff/mothers who may be pregnant) Tonsillitis None once child is well	Measles	•	
Meningococcal Meningitis/Septicaemia Will give specific advice on any action needed. Siblings need not be excluded Meningitis None once child is well infection risk is minimal Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea 48 hours after the last bout of either. Slapped Cheek Syndrome (Parvovirus B19) (Kool Kids must notify staff/mothers who may be pregnant) Tonsillitis None once child is well		1 '	
Meningitis/Septicaemiawill give specific advice on any action needed. Siblings need not be excludedMeningitisNone once child is well infection risk is minimalMumpsExclude child for five days after onset of swellingScabiesUntil treatedScarlet FeverChild can return 24 hours after commencing appropriate antibiotic treatmentSickness & diarrhoea48 hours after the last bout of either.Slapped Cheek Syndrome (Parvovirus None.None.B19)(Kool Klds must notify staff/mothers who may be pregnant)TonsillitisNone once child is well			
Siblings need not be excluded Meningitis None once child is well infection risk is minimal Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea 48 hours after the last bout of either. Slapped Cheek Syndrome (Parvovirus None. B19) (Kool Klds must notify staff/mothers who may be pregnant) Tonsillitis None once child is well			
MeningitisNone once child is well infection risk is minimalMumpsExclude child for five days after onset of swellingScabiesUntil treatedScarlet FeverChild can return 24 hours after commencing appropriate antibiotic treatmentSickness & diarrhoea48 hours after the last bout of either.Slapped Cheek Syndrome (Parvovirus B19)None. (Kool Klds must notify staff/mothers who may be pregnant)TonsillitisNone once child is well	Meningitis/Septicaemia	,	
Mumps Exclude child for five days after onset of swelling Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea 48 hours after the last bout of either. Slapped Cheek Syndrome (Parvovirus None. B19) (Kool Klds must notify staff/mothers who may be pregnant) Tonsillitis None once child is well	Moningitie		
Scabies Until treated Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea 48 hours after the last bout of either. Slapped Cheek Syndrome (Parvovirus None. B19) (Kool Klds must notify staff/mothers who may be pregnant) Tonsillitis None once child is well			
Scabies Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea Slapped Cheek Syndrome (Parvovirus B19) Fifth Disease Tonsillitis Until treated Child can return 24 hours after commencing appropriate antibiotic treatment None appropriate antibiotic treatment None. (Kool Klds must notify staff/mothers who may be pregnant) None once child is well	wiumps	•	
Scarlet Fever Child can return 24 hours after commencing appropriate antibiotic treatment Sickness & diarrhoea 48 hours after the last bout of either. Slapped Cheek Syndrome (Parvovirus None. (Kool Klds must notify staff/mothers who may be pregnant) Tonsillitis None once child is well	Scapios		
appropriate antibiotic treatment Sickness & diarrhoea 48 hours after the last bout of either. Slapped Cheek Syndrome (Parvovirus None. (Kool Klds must notify staff/mothers who may be Fifth Disease pregnant) Tonsillitis None once child is well			
Sickness & diarrhoea 48 hours after the last bout of either. Slapped Cheek Syndrome (Parvovirus None. (Kool Klds must notify staff/mothers who may be pregnant) Tonsillitis None once child is well	Scarlet i ever	_	
Slapped Cheek Syndrome (Parvovirus B19) Fifth Disease Tonsillitis None. (Kool Klds must notify staff/mothers who may be pregnant) None once child is well	Sickness & diarrhoea	1	
B19) (Kool Klds must notify staff/mothers who may be pregnant) Tonsillitis None once child is well			
Fifth Disease pregnant) Tonsillitis None once child is well	, , ,		
Tonsillitis None once child is well	•	1 '	
VINCODING COUGH J MAVO HOHI COHHILICHICHICH OF HEALHICHE	Whooping Cough	5 days from commencement of treatment	

This list is statutory guidance from the NHS and is not negotiable. The setting reserves the right to turn away any family who breaches this policy.

This policy was adopted at a meeting of	Kool Kids Club	
Held in:	April 2024	
To be reviewed in:	April 2025	
Signed on behalf of the setting:		
Name of Signatory:		
Role of Signatory:	Manager	Deputy Manager